



## Complete Summary

---

### TITLE

Advanced chronic kidney disease (CKD): percent of patients referred for surgery for construction of an arteriovenous (AV) fistula on index date.

### SOURCE(S)

Renal Physicians Association. Appropriate patient preparation for renal replacement therapy. Rockville (MD): Renal Physicians Association; 2002 Oct 1. 78 p. (Clinical Practice Guideline; no. 3).

## Brief Abstract

### DESCRIPTION

This measure assesses the percent of patients referred for surgery for construction of an arteriovenous (AV) fistula on index date (i.e., the date patient is first seen by practitioner and identified as having advanced chronic kidney disease [CKD]) among patients with advanced CKD for whom hemodialysis is the chosen mode of renal replacement therapy.

### RATIONALE

Utilizing conventional criteria for the initiation of renal replacement therapy (RRT), such as congestive heart failure, malnutrition, acidosis or uremia, can lead to patients being treated only when they have already suffered irreversible cumulative complications from chronic kidney disease (CKD). It is therefore imperative to properly time the initiation of RRT in order to minimize morbidity and mortality. The role of patient factors such as therapeutic preferences is also considered especially important in timing, initiation, and choice of modality of RRT.

Patients and healthcare professionals should be educated about the need to preserve veins to avoid loss of potential access sites. Repeated venipuncture for vascular access may render arm vein sites unsuitable for construction of a primary arteriovenous (AV) fistula. In addition, subclavian vein catheterization should be avoided for temporary access due to the risk of central venous stenosis.

### PRIMARY CLINICAL COMPONENT

Advanced chronic kidney disease; arteriovenous fistula; surgery

### DENOMINATOR DESCRIPTION

The number of adult patients with advanced chronic kidney disease (CKD) not currently receiving renal replacement therapy (RRT) for whom hemodialysis is the chosen mode of RRT

#### NUMERATOR DESCRIPTION

The number of patients from the denominator referred for surgery for construction of an arteriovenous (AV) fistula on index date (i.e., the date patient is first seen by practitioner and identified as having advanced advanced chronic kidney disease [CKD])

#### Evidence Supporting the Measure

##### PRIMARY MEASURE DOMAIN

Process

##### SECONDARY MEASURE DOMAIN

Not applicable

##### EVIDENCE SUPPORTING THE MEASURE

A clinical practice guideline or other peer-reviewed synthesis of the clinical evidence

A formal consensus procedure involving experts in relevant clinical, methodological, and organizational sciences

A systematic review of the clinical literature

##### NATIONAL GUIDELINE CLEARINGHOUSE LINK

- [Appropriate patient preparation for renal replacement therapy.](#)

#### Evidence Supporting Need for the Measure

##### NEED FOR THE MEASURE

Unspecified

#### State of Use of the Measure

##### STATE OF USE

Pilot testing

##### CURRENT USE

Internal quality improvement

## Application of Measure in its Current Use

### CARE SETTING

Ambulatory Care  
Physician Group Practices/Clinics

### PROFESSIONALS RESPONSIBLE FOR HEALTH CARE

Physicians

### LOWEST LEVEL OF HEALTH CARE DELIVERY ADDRESSED

Individual Clinicians

### TARGET POPULATION AGE

Age greater than or equal to 18 years

### TARGET POPULATION GENDER

Either male or female

### STRATIFICATION BY VULNERABLE POPULATIONS

Unspecified

## Characteristics of the Primary Clinical Component

### INCIDENCE/PREVALENCE

Unspecified

### ASSOCIATION WITH VULNERABLE POPULATIONS

Unspecified

### BURDEN OF ILLNESS

Unspecified

### UTILIZATION

Unspecified

### COSTS

Unspecified

## Institute of Medicine National Healthcare Quality Report Categories

### IOM CARE NEED

Living with Illness

### IOM DOMAIN

Effectiveness

Patient-centeredness

## Data Collection for the Measure

### CASE FINDING

Users of care only

### DESCRIPTION OF CASE FINDING

Adult patients 18 years and older with advanced chronic kidney disease (CKD) for whom hemodialysis is the chosen mode of renal replacement therapy (RRT)

### DENOMINATOR (INDEX) EVENT

Clinical Condition

Therapeutic Intervention

### DENOMINATOR INCLUSIONS/EXCLUSIONS

#### Inclusions

Adult patients age 18 years and older with chronic kidney disease stage 4 or 5 (glomerular filtration rate [GFR] less than or equal to 30 mL/min/1.73 m<sup>2</sup>), not currently receiving renal replacement therapy, for whom hemodialysis is the chosen mode of renal replacement therapy (RRT)

#### Exclusions

Unspecified

### NUMERATOR INCLUSIONS/EXCLUSIONS

#### Inclusions

The number of patients from the denominator referred for surgery for construction of an arteriovenous (AV) fistula on index date (i.e., the date patient is first seen by practitioner and identified as having advanced chronic kidney disease [CKD])

#### Exclusions

Unspecified

### DENOMINATOR TIME WINDOW

Time window precedes index event

#### NUMERATOR TIME WINDOW

Episode of care

#### DATA SOURCE

Administrative data  
Laboratory data  
Medical record

#### LEVEL OF DETERMINATION OF QUALITY

Individual Case

#### PRE-EXISTING INSTRUMENT USED

Unspecified

### Computation of the Measure

#### SCORING

Rate

#### INTERPRETATION OF SCORE

Better quality is associated with a higher score

#### ALLOWANCE FOR PATIENT FACTORS

Unspecified

#### STANDARD OF COMPARISON

Internal time comparison

### Evaluation of Measure Properties

#### EXTENT OF MEASURE TESTING

Unspecified

### Identifying Information

#### ORIGINAL TITLE

Number of patients referred for surgery for construction of an AV fistula on index date / number of patients with advanced CKD for whom hemodialysis is the chosen mode of RRT.

#### MEASURE COLLECTION

[Renal Physicians Association Clinical Performance Measures on Appropriate Patient Preparation for Renal Replacement Therapy](#)

#### MEASURE SET NAME

[Renal Physicians Association Clinical Performance Measures for Timing of Renal Replacement Therapy Recommendations](#)

#### DEVELOPER

Renal Physicians Association

#### ADAPTATION

Measure was not adapted from another source.

#### RELEASE DATE

2002 Oct

#### MEASURE STATUS

This is the current release of the measure.

#### SOURCE(S)

Renal Physicians Association. Appropriate patient preparation for renal replacement therapy. Rockville (MD): Renal Physicians Association; 2002 Oct 1. 78 p. (Clinical Practice Guideline; no. 3).

#### MEASURE AVAILABILITY

The individual measure, "Number of patients referred for surgery for construction of an AV fistula on index date / number of patients with advanced CKD for whom hemodialysis is the chosen mode of RRT," is published in "Renal Physicians Association Clinical Practice Guideline #3: Appropriate Patient Preparation for Renal Replacement Therapy."

For more information, contact RPA at 1700 Rockville Pike, Suite 220, Rockville, MD 20852; phone: 301-468-3515; fax: 301-468-3511; Web site: [www.renalmd.org](http://www.renalmd.org); e-mail: [rpa@renalmd.org](mailto:rpa@renalmd.org).

#### NQMC STATUS

This NQMC summary was completed by ECRI on May 23, 2003. The information was verified by the Renal Physicians Association on June 17, 2003.

#### COPYRIGHT STATEMENT

This NQMC summary is based on the original measure, which is subject to the measure developer's copyright restrictions.

For more information, contact RPA at 1700 Rockville Pike, Suite 220, Rockville, MD 20852; phone: 301-468-3515; fax: 301-468-3511; Web site: [www.renalmd.org](http://www.renalmd.org); e-mail: [rpa@renalmd.org](mailto:rpa@renalmd.org).

© 2004 National Quality Measures Clearinghouse

Date Modified: 10/25/2004

**FIRSTGOV**

